

1) The arterial supply of the corpus cavernosum is usually from the:

- **a** external pudendal artery.
- b** accessory pudendal artery.
- c** cavernous artery.
- d** the dorsal artery.
- e** inferior epigastric artery

2) Reduced penile venous outflow during erection is due to:

- **a** active constriction of the superficial and deep dorsal veins.
- b** opening of penile arterio-venous shunts.
- c** compression of the subtunical venules and emissary veins by the tunica albuginea.
- d** active constriction of emissary veins.
- e** relaxation of the ischiocavernosus muscle

3) The innervation of the penis comes from:

- **a** S2-4 spinal segments.
- b** T10-12 and S2-4 spinal segments.
- c** the dorsal nerve.
- d** the cavernous nerve.
- e** all of the above.

4) The principal neurotransmitter mediating penile erection is:

- **a** prostaglandin E 1.
- b** nitric oxide.
- c** norepinephrine.
- d** acetylcholine.
- e** neuropeptide P.

5) ED associated with diabetes can be a result of:

- **a** psychologic impact.
- b** neurologic deficit.
- c** arterial insufficiency.
- d** endothelial cell dysfunction.
- e** all of the above.

- 6) A patient with ED who has a low risk for cardiovascular events should:
- **a** undergo cardiac stress testing.
 - b** receive cardiologic referral.
 - c** receive counseling for lifestyle modifications
 - d** initiate statin therapy.
 - e** be reassured that there is no need for cardiovascular health intervention.

7) Recommended serum laboratory testing for men with sexual problems typically include:

- **a** a hemoglobin A1C level.
- b** a lipid profile.
- c** thyroid function tests.
- d** a prolactin level.
- e** a PSA measurement.

8) The best indicator of hypogonadism is:

- **a** a screening questionnaire.
- b** a total serum testosterone measurement.
- c** a free serum testosterone measurement.
- d** a bioavailable serum testosterone measurement.
- e** an intratesticular testosterone measurement.

9) Optimization of effect for PDE5 inhibitors is best achieved by:

- **a** daily dosing.
- b** reducing the frequency of sexual intercourse attempts.
- c** increasing food intake with dosing.
- d** applying sexual stimulation.
- e** using twice-maximal dosages.

10) Infected penile prostheses are best treated by:

- **a** removal of the single infected component.
- b** removal of all prosthetic components.
- c** twelve weeks of broad spectrum antibiotics.
- d** hyperbaric oxygen.
- e** 12 weeks of broad-spectrum antibiotics + hyperbaric oxygen

1	C
2	C
3	E
4	B
5	E
6	C
7	B
8	D
9	D
10	B